

# Pourakino Catchment Group

Fresh water health and landscape influences  
in Pourakino Catchment



**THRIVING  
SOUTHLAND**

*Tōtini ana te whenua. Tōtini ana te takata.  
A thriving, prosperous land. A thriving, prosperous people.*

AS AT NOVEMBER 2021  
Further updates will be included as new  
information becomes available.

# Welcome to Pourakino Catchment brochure

This Pourakino Catchment Group brochure is one of a series of brochures on catchments in Southland.

Catchment Groups have been asking for more detailed insights into their individual catchments. This brochure provides insights based on available information, bringing together published science, research, data and information on the state of water, soil and land in Pourakino Catchment.

It provides details on what affects water quality and how these impact the rest of the catchment, including other catchment areas that may be impacted by what goes on in the catchment, such as the estuary.

Although this brochure collates all the available information that has been brought together in a literature review commissioned by Thriving Southland - the Science Report [thrivingsouthland.co.nz/science-report](https://thrivingsouthland.co.nz/science-report) - it may not have all the details you know about in your catchment, or the research you may have done on your farm or in your Catchment Group.

## How to use this brochure

This brochure sets out (publicly available) details on:

- » Water quality
- » Landscape influences
- » Physiographic zones
- » Groundwater management zones (GMZs)
- » Measuring what lives in streams and rivers
- » Macroinvertebrate community index (MCI)
- » Estuary health
- » Where to get more information.

If you are not familiar with the terms and language used, read the brochure in conjunction with this glossary [environment.govt.nz/publications/environmentaotearoa-2019-glossary](https://environment.govt.nz/publications/environmentaotearoa-2019-glossary).

We also recommend you check out the Catchment Group page on <https://thrivingsouthland/pourakino/> to learn more about the catchment and what projects the Catchment Group has underway or planned.

## Interpreting what the data in the brochure means

Because this brochure brings together the data available, we have deliberately not interpreted that data or explained what the trends may mean for your catchment.

We recommend you contact an environmental consultant, your Thriving Southland Catchment Group coordinator or Environment Southland to speak to experts who can explain what these trends and data may mean for your catchment, or for your farm specifically.

You can also check out a range of information on the Thriving Southland Information Resource Hub [thrivingsouthland.co.nz/infohub](https://thrivingsouthland.co.nz/infohub) which will connect you with tools and resources from many different organisations to help with understanding limit setting, environmental contributing factors, mitigations and options available to you.

## A little bit about Thriving Southland

Thriving Southland supports Southland's Catchment Groups to understand challenges and opportunities in their catchments and create innovative and exciting solutions.

We have a vision to create a prosperous Southland, healthy people, and a healthy environment, and believe that by working together, Thriving Southland's communities will create a better future for all by protecting the region's prosperity, heritage, environment and health.

## Thanks

Thank you to the farmers who supported the development of this brochure, and to the Ministry of Primary Industry for its Sustainable Land Use Programme which supports the work Thriving Southland is delivering for farmers and communities in Southland. Thank you also to Environment Southland who reviewed the content of this brochure.



# Water quality in Aparima Catchment

## Aparima Catchment

Pourakino is part of the Aparima Catchment which outflows into the Jacobs River Estuary. The Jacobs River Estuary are an important source of mahinga kai, particularly shellfish, mussels, paua, tuna and inanga.

Currently the Jacobs River Estuary is considered to be in fair to poor condition.

## Summary of Pourakino Catchment


The hydrology, health and functions of a stream/river or area of groundwater are directly linked to the characteristics of its catchment, including the landscape, soils and human activities.

- » This catchment covers a mosaic of four different physiographic zones that differ in nitrate and phosphorus levels from land use.
- » Part of the Pourakino catchment area overlies part of the Lower Aparima GMZ, which can have elevated nitrate levels in shallow limestone aquifers.
- » Water quality in this catchment is generally good but is showing stress in terms of *E. coli* (faecal bacteria) (surface water), and nitrogen in some areas (groundwater and surface water).
- » Neighbouring properties on different zones may have very different water quality outcomes with similar land practices, due to different contamination movement and attenuation pathways (reducing the effects of contaminants).



## What does this mean?

- » In August 2020, the Government announced changes to the National Policy Statement for Freshwater Management (NPS-FM), the National Environmental Standard for Freshwater (NES) and changes to the Resource Management Act. These outline changes in regulations relating to wetlands and rivers, intensive winter grazing, intensification, stockholding areas and stock exclusion.
- » More about Environment Southland's response to the Government's Essential Freshwater Package [es.govt.nz/environment/water/essential-freshwater-package](https://es.govt.nz/environment/water/essential-freshwater-package)



- » Environment Southland, in partnership with Te Ao Mārama Inc\*, has indicated a proposed limits and targets plan change (LTPC) will be notified in 2023. This will establish nutrient limits and targets to improve the quality of groundwater and surface water. [waterandland.es.govt.nz/about/values-and-objectives](https://waterandland.es.govt.nz/about/values-and-objectives)
- » Environment Southland and Te Ao Mārama Inc have established a community-based regional forum to consider and advise on limits, targets, and methods.
- » Plan changes will result in additional controls and rules in Southland that will be focused on reducing the loss of nutrients, specifically nitrogen and phosphorus, and reducing discharges of sediment and faecal microorganisms, from land to groundwater and surface water.
- » In the Environment Southland Proposed Water and Land Plan there is a focus on good management practices (GMPs) and farm environmental management plans (FEPs). You can view GMP factsheets for each physiographic zone on the Environment Southland website [es.govt.nz](https://es.govt.nz)

\* Te Ao Mārama Incorporated looks after mana whenua interests in resource management and other aspects related to local government in Southland. It is authorised to represent three Ngāi Tahu papatipu runanga in Murihiku/Southland. It is involved in the protection of the spiritual and cultural values of the region, including wahi tapu (sacred places), mahinga kai (gathering of food and resources) and other natural resources.

# Pourakino water quality

Surface water quality is assessed by testing how much nitrogen, phosphorus and *E. coli* is present. LAWA summary results for this catchment are shown below ([lawa.org.nz](http://lawa.org.nz)):

## Total oxidised nitrogen

Monitoring site	5-year median	5-year trend	10-year trend	15-year trend
Cascade Stream at Pourakino Valley Rd	0.02 mg/L			
Opouriki Stream at Tweedie Rd	1.97 mg/L			
Pourakino River at Traill Road	0.197 mg/L			

<sup>^</sup> TON is the sum of nitrate and nitrite. Nitrite is generally a very small fraction of the TON concentration in rivers, TON is taken to be equivalent to the nitrate concentration

\* 2016-2020 LAWA median per NPS-FM 2020 using TON as surrogate for N03-N

\* Too much TON can contribute to excessive algal growth in waterways.

## Ammoniacal nitrogen

Monitoring site	5-year median	State	5-year trend	10-year trend	15-year trend
Cascade Stream at Pourakino Valley Rd	0.005 mg/L				
Opouriki Stream at Tweedie Rd	0.021 mg/L				
Pourakino River at Traill Road	0.012 mg/L				

\* If ammoniacal nitrogen reaches very high concentrations it can become toxic under certain temperature and pH conditions.

## Dissolved reactive phosphorus

Monitoring site	5-year median	State	5-year trend	10-year trend	15-year trend
Cascade Stream at Pourakino Valley Rd	0.002 mg/L				
Opouriki Stream at Tweedie Rd	0.009 mg/L				
Pourakino River at Traill Road	0.002 mg/L				

\* Dissolved reactive phosphorus concentrations are an indication of a waterbody's ability to support nuisance algal or plant growths (algal blooms).

## Total phosphorus

Monitoring site	5-year median	5-year trend	10-year trend	15-year trend
Cascade Stream at Pourakino Valley Rd	0.008 mg/L			
Opouriki Stream at Tweedie Rd	0.035 mg/L			
Pourakino River at Traill Road	0.013 mg/L			

\* Too much phosphorus can encourage the growth of nuisance plants such as algal blooms.

KEY (STATE)		KEY (TREND)		
 <b>A</b> Very good	 <b>B</b> Good	 <b>Very likely improving</b>	 <b>Likely improving</b>	 <b>Indeterminate</b>
 <b>C</b> Fair	 <b>D</b> Poor	 <b>Very Likely degrading</b>	 <b>Likely Degrading</b>	 <b>Not Assessed</b>

## E. coli

Monitoring site	5-year median	State	5-year trend	10-year trend	15-year trend
Cascade Stream at Pourakino Valley Rd	110n/100 mL	 <b>A</b>			
Opouriki Stream at Tweedie Rd	575n/100 mL	 <b>E</b>			
Pourakino River at Traill Road	345n/100 mL	 <b>E</b>			

\* High concentrations of this bacteria exceeding water quality guidelines indicate faecal contamination which can be harmful to humans.

\* 2016-2020 LAWA median graded as per NPS-FM 2020

KEY (STATE)			KEY (TREND)		
 <b>A</b> Very good (infection risk is 1%)	 <b>B</b> Good (infection risk is 2%)	 <b>C</b> Fair (infection risk is 3%)	 <b>Very likely improving</b>	 <b>Likely improving</b>	 <b>Indeterminate</b>
 <b>D</b> Poor (infection risk is >3%)	 <b>E</b> Very Poor (infection risk is >7%)		 <b>Very Likely degrading</b>	 <b>Likely Degrading</b>	 <b>Not Assessed</b>

Results from lawa.org.nz (October 2021)

# MCI

Macroinvertebrates include the caddisflies, mayflies, stoneflies, worms and snails that live in rivers. They are an important food source for fish and birds and sensitive to the combination of nutrients, sediment and habitat. Because of this sensitivity, they are considered to be a good representation of overall water quality and ecosystem health. The different macroinvertebrates present can be identified and then converted to a score called the Macroinvertebrate Community Index (MCI).

A higher MCI score generally indicates a healthier stream. Generally, MCI scores range from >150 (very good water quality) to as low as 20 (very poor water quality).

Results for this catchment are (LAWA October 2021):

## MCI

Monitoring site	5-year median	State	10-year trend	15-year trend
Cascade Stream at Pourakino Valley Rd	118.0			
Opouriki Stream at Tweedie Rd	NO DATA	NO DATA	NO DATA	NO DATA
Pourakino River at Traill Road	NO DATA	NO DATA	NO DATA	NO DATA

### KEY (STATE)

A

**Macroinvertebrate community indicative of pristine conditions with almost no organic pollution or nutrient enrichment.**

B

**Macroinvertebrate community indicative of mild organic pollution or nutrient enrichment. Largely composed of taxa sensitive to organic pollution/nutrient enrichment.**

C

**Macroinvertebrate community indicative of moderate organic pollution or nutrient enrichment. There is a mix of taxa sensitive and insensitive to organic pollution/nutrient enrichment.**

D

**Macroinvertebrate community indicative of severe organic pollution or nutrient enrichment. Communities are largely composed of taxa insensitive to inorganic pollution/nutrient enrichment.**

National bottom line: MCI score 90

### KEY (TREND)

Very likely improving

Likely improving

Indeterminate

Very Likely degrading

Likely Degrading

Not Assessed

# Estuary health

Table: Estuary state information (provided by Environment Southland July 2021, es.govt.nz state and outcome map).

The Pourakino River flows into the Jacobs River Estuary and is an important factor in understanding the impacts of water quality in Pourakino. Decisions made in Pourakino affect water quality upstream, flow downstream and impact on water quality in the estuary.



The below assessment of estuary health based on the sediment quality gives a good indication of what is happening upstream across all the catchments that feed into the waterways supplying the estuary and therefore impact on the ecosystems and the cultural values of the area.

Estuary	Soft mud	Nutrients in sediment	Oxygen in sediment	Macroalgae cover	Seagrass loss	GEZ*
Waiau Lagoon/Te Wae		Orange	Yellow			
Lake Brunton			Green			
Waituna Lagoon/Waiparera (not assessed)						
New River Estuary	Orange	Orange	Orange	Yellow	Orange	Orange
Jacobs River Estuary	Orange	Yellow	Orange	Orange	Orange	Orange
Waikawa Estuary	Orange	Yellow	Orange	Green	Orange	Green
Haldane Estuary	Yellow	Green	Teal	Teal		Teal
Freshwater Estuary	Teal	Green	Teal	Green	Yellow	Teal
Waimatuku Estuary		Yellow	Yellow			
Toetoes Estuary	Yellow	Green	Orange	Yellow	Orange	Green

**KEY** Very Good Good Fair Poor

\* Gross Eutrophic Zone is defined as an area that has low sediment oxygenation (<1cm aRPD), soft mud (>25% mud content) and the presence of high macroalgal cover (>50% cover), these areas are in poor condition and can no longer support most estuarine animals and shellfish.

## Aparima Catchment with Estuaries and Surface water quality monitoring sites\*



\*Sites in Pourakino Catchment only shown

# Landscape influences

What we do on the land can affect our water, but how it affects our water depends on a range of factors, including how our landscape works. It is useful to look at:

- » Physiographic zones which help to explain how nitrogen, phosphorus, sediment and faecal microorganisms (such as *E. coli*) move and are attenuated (reduced, e.g. by biological or chemical processes)
- » GMZs which highlight the connectivity between land, surface water and groundwater.

## **Physiographic zones**

Southland has been divided into nine physiographic zones to help understand how water moves across the landscape and why water quality is better in some places than others. Each physiographic zone represents an area that has similar factors influencing water quality, such as climate, topography, geology and soil type.

Most of the Pourakino Catchment area falls into the bedrock/hill country physiographic zone. To the south, the catchment area is dominated by the lignite-marine terraces physiographic zone. Areas of oxidising and gleyed physiographic zones are also found (see map below). These zones differ in the way contaminants are transported and attenuated within the catchment.



# Pourakino Catchment showing physiographic zones

## Bedrock/Hill Country – overland flow

This zone is found on rolling to steep land below 800 metres. It is characterized by high rainfall and a dense network of branching streams.

Water quickly flows down-slope to nearby streams following high or prolonged rainfall. Nitrogen, phosphorus, sediment and faecal microorganisms are all carried with water, particularly during late autumn and winter.

## Lignite-marine terraces - overland flow

Located on gently undulating to rolling land that have a high potential for overland flow.

## Lignite-marine terraces - artificial drainage

Located on gently undulating land that has slow subsoil permeability and may be seasonally wet.

Organic-rich sediments like lignite, are found at depth within this zone. The presence of these sediments has a strong influence on reducing the amount of nitrate in groundwater. Denitrification\* rates are high.

## Bedrock/Hill country - artificial drainage

Generally located on developed land along the base of hillslopes. Artificial drainage is needed in areas of low slope and low subsoil permeability.

## Oxidising - overland flow

The overland flow variant is found on steeper areas where water preferentially flows over the land surface.

## Oxidising

Soils and aquifers in this zone have high risk of nitrogen build-up due to low rates of denitrification\*

The combination of flat land and well drained soils results in high rates of nitrogen leaching (deep drainage) to underlying aquifers.

## Gleyed

This zone is generally found in areas that were once wetlands. It is characterized by a dense network of streams and a high water-table during winter.


Soils are prone to waterlogging and have some denitrification\* ability, which reduces build-up of soil nitrogen. However, an extensive network of artificial drainage rapidly transport nitrogen, phosphorus, sediment and faecal microbes to surface water, particularly during heavy rain.



\*Denitrification occurs when nitrate is converted to nitrogen gas via various reactions involving bacteria. Where denitrification occurs, nitrogen is effectively 'lost' from soil and water as a gas. This is a form of attenuation.

# GMZ - Pourakino


Parts of the eastern side of the Pourakino Catchment area overlies part of the Lower Aparima GMZ.



# Lower Aparima GMZ

The Lower Aparima GMZ covers approximately 29,000ha in the lower reaches of the Aparima River Catchment:

- » Depth to groundwater ranges from less than 2 metres below ground level on Aparima River floodplain to 20 metres below ground level in limestone aquifers underlying higher terraces.
- » Seasonal groundwater variation is generally less than 2 metres, but can be up to 10 metres in limestone aquifers.
- » A diagrammatic cross-section of this GMZ showing areas of groundwater is provided below (source [es.govt.nz/environment/water/groundwater/groundwater-management-zones/lower-aparima](https://es.govt.nz/environment/water/groundwater/groundwater-management-zones/lower-aparima)).
- » Groundwater recharge in this zone is derived from local rainfall and runoff from surrounding hills, which soaks through the soil. Groundwater discharge mostly occurs as springs and as baseflow into Aparima River.



## Quaternary sediments

- Q1 Alluvium
- Q2 Alluvium
- Alluvial Fan
- Q6-8 Alluvium

## Tertiary sediments

- Winton Hill Formation mudstone
- Winton Hill Formation sandstone
- Winton Hill Formation lignite
- Forest Hill Formation limestone

## Basement rock

- Brook Street Terrane (volcaniclastic sediments)

## Groundwater quality Lower Aparima GMZ

- » Nitrate = generally low but may be elevated in shallow limestone aquifers
- » Phosphorus = low
- » *E. coli* = low, but risk may be elevated close to source.



## Find out more

**Environment Southland Aparima flood warning fact sheet**

[bit.ly/3cLate5](https://bit.ly/3cLate5)

**Find out more about physiographic zones**

[bit.ly/2OI7z7F](https://bit.ly/2OI7z7F)

**Find out more about Southland's groundwater**

[bit.ly/30Db5g1](https://bit.ly/30Db5g1)

---

## Find out more about stream health

**Environment Southland**

[es.govt.nz/environment/water/rivers-and-streams](https://es.govt.nz/environment/water/rivers-and-streams)

**Land Air Water Aotearoa (LAWA)**

[lawa.org.nz](https://lawa.org.nz)

**Ministry for the Environment**

[environment.govt.nz/facts-and-science/freshwater](https://environment.govt.nz/facts-and-science/freshwater)

**Link to iwi freshwater objectives**

[bit.ly/2P4HsBV](https://bit.ly/2P4HsBV)

---

## Get in contact

**For more information about your catchment and to contact your local catchment coordinator**

021 466 700 | [office@thrivingsouthland.co.nz](mailto:office@thrivingsouthland.co.nz)

[thrivingsouthland.co.nz/catchment-groups](https://thrivingsouthland.co.nz/catchment-groups)



# THRIVING SOUTHLAND

*Tōnui ana te whenua. Tōnui ana te takata.  
A thriving, prosperous land. A thriving, prosperous people.*